

Date: <u>9/23/98</u>	Express Mail Label No. <u>EM012016453US</u>
----------------------	---

Inventor: Francis A. Ennis
Attorney's Docket No.: UMMC98-13

PREDICTIVE ASSAY FOR IMMUNE RESPONSE

BACKGROUND OF THE INVENTION

Among the greatest successes in the field of public health is widespread vaccination against a variety of formerly common infectious diseases. For example,
5 public vaccination programs in the United States have eradicated smallpox and dramatically reduced the incidence of diseases such as measles, rubella, polio and diphtheria, among others. However, the development of novel vaccine compositions is still an active area of research. In particular, the development of effective vaccines for a number of diseases for which no clinically proven vaccine exists remains an important
10 goal. For example, a vaccine which protects against infection by human immunodeficiency virus (HIV) is a primary goal in efforts to control the spread of AIDS. Also needed are vaccine compositions which have improved efficacy in comparison to vaccines in current use.

The efficacy of a vaccine for use in humans depends upon the ability of the
15 vaccine formulation to elicit an immune response which is sufficient to provide protection against subsequent challenge with the pathogen. Experimental vaccines are typically evaluated first *in vivo* in small animals, such as mice, guinea pigs or rabbits. The assessment of the experimental vaccines generally relies upon measurements of serum antibody responses and, sometimes, antigen-specific lymphocyte proliferative
20 responses. Vaccine formulations which are successful in these animal models are then tested in sub-human primates and, finally, in humans.

The assessment of a test vaccine in an animal model is costly and takes considerable time. Typically, several doses of vaccine are administered to the animal at

intervals of several weeks. The immune response of primates to a given test vaccine is often less than that of smaller animals, and clinical studies in humans are ultimately required to determine the efficacy of a test vaccine. In addition to the large costs associated with purchasing and housing animals for long periods of time, each step of the process requires a minimum of several months. Thus, the number of experimental vaccines which can be evaluated using prior art methods is necessarily limited, with the possible result that potentially useful vaccine formulations may never be tested.

There is, therefore, a need for an *in vitro* test for determining the human immune response to an experimental vaccine construct which would allow the rapid evaluation of large numbers of candidate vaccine compositions within a short time period and at reasonable cost.

SUMMARY OF THE INVENTION

The present invention relates to a method for assessing the ability of a candidate vaccine composition to stimulate a T cell response. In one embodiment, the invention provides a method for selecting one or more vaccine compositions from among a group of vaccine compositions for *in vivo* assessment, for example, in one or more animal or human subjects. Each of the vaccine compositions comprises one or more antigens or one or more nucleic acid molecules encoding one or more antigens. The method comprises the steps of: (1) contacting antigen presenting cells in culture with a vaccine composition selected from among the group of vaccine compositions, thereby, if one or more of the antigens or nucleic acid molecules can be taken up and processed by the antigen presenting cells, producing one or more processed antigens; (2) contacting the antigen presenting cells with T cells under conditions sufficient for the T cells to respond to one or more of the processed antigens; (3) determining whether the T cells respond to one or more of the processed antigens; whereby if the T cells respond to one or more of the processed antigens, then the vaccine composition stimulates a T cell response; and (4) repeating steps (1), (2) and (3) with each additional vaccine composition in the group, thereby identifying the vaccine compositions within the group which stimulate a T cell response; and, if one or more of these vaccine compositions stimulates a T cell response, (5) selecting at least one vaccine composition which

stimulates a T cell response for assessment in one or more animals and/or in one or more human subjects.

In another embodiment, the invention relates to a method of selecting a vaccine composition from a group consisting of two or more vaccine compositions for
5 assessment in one or more animals or in one or more human subjects. Each of the vaccine compositions comprises one or more antigens or one or more nucleic acid molecules encoding one or more antigens. The method comprises the steps of:

- (1) contacting antigen presenting cells in culture with a vaccine composition selected from among said group of vaccine compositions, thereby, if one or more of the antigens
10 or nucleic acid molecules are taken up and processed by the antigen presenting cells, producing one or more processed antigens; (2) contacting the antigen presenting cells with T cells under conditions sufficient to produce a T cell response to one or more of the processed antigens, thereby producing a vaccine composition-stimulated T cell response; (3) measuring the vaccine composition-stimulated T cell response;
15 (4) repeating steps (1), (2) and (3) with each of the remaining vaccine compositions in the group, thereby identifying the vaccine composition or compositions which stimulate the greatest T cell response; (5) selecting the vaccine composition or compositions which stimulate the greatest T cell response for assessment in one or more animals and/or in one or more human subjects.

- 20 In a further embodiment, the invention relates to a method for assessing the ability of a vaccine composition comprising one or more antigens or one or more nucleic acid molecules encoding one or more antigens to stimulate a protective T cell response. The method comprises the steps of: (1) contacting human antigen presenting cells in culture with the vaccine composition, thereby, if one or more of the antigens or
25 nucleic acid molecules can be taken up and processed by the antigen presenting cells, producing one or more processed antigens; (2) contacting the antigen presenting cells with human T cells under conditions sufficient to produce a T cell response to one or more of the processed antigens, thereby producing a T cell response; (3) measuring the T cell response; and, if the T cell response is greater than a pre-selected value,
30 (4) assessing the ability of the vaccine composition to stimulate a protective T cell response in one or more animals or in one or more human subjects.

In another embodiment, the method of the invention comprises the steps of :

- (1) contacting human antigen presenting cells in culture with the vaccine composition, whereby, if one or more of the antigens are taken up and processed by the antigen presenting cells, said antigen or antigens are processed by the antigen presenting cells, thereby producing one or more processed antigens; (2) contacting the antigen presenting cells of step (1) with human T cell clones which are specific for an epitope within one or more of the antigens for a period of time sufficient for the human T cell clones to respond to one or more of the processed antigens; and (3) determining whether the human T cell clones respond to the processed antigen or antigens. If the T cell clones respond to the processed antigen or antigens, the method can, optionally, further include the step of assessing the vaccine composition in one or more animals or human subjects.

- Preferably, the vaccine composition includes at least one antigen which comprises a T cell epitope, and the T cells are T cell clones which are specific for a T cell epitope in at least one of the antigens. In one embodiment, the T cells are CD8⁺ T cells and the vaccine composition includes at least one antigen comprising antigen a CD8 epitope. In this embodiment, the T cell response to the processed antigen can be, for example, T cell proliferation, cytolysis of the antigen presenting cells or the production of one or more cytokines.

- In another embodiment, the T cells are CD4⁺ T cells and the vaccine composition includes at least one antigen which comprises a CD4 epitope. In this embodiment, the T cell response to the processed antigen which is determined can be, for example, T cell proliferation, stimulation of antibody production by B cells or production of one or more cytokines.

- The present invention offers several advantages over prior art methods of evaluating candidate vaccine compositions. For example, the method of the invention can be completed in a relatively short time period. The present method can also be used as a first screen to determine which candidate compositions should be evaluated in much more expensive and time consuming *in vivo* tests. Thus, the method of the invention enables the efficient and cost effective evaluation of large numbers of potential vaccine compositions, increasing the possibility that effective vaccine compositions will be discovered.

Cell-mediated immunity is dependent upon lymphocytes known as B cells and T cells. B cells produce antibodies targeted against extracellular antigens. T cells recognize antigen fragments (peptides) which are displayed at the surface of a host cell. Such antigen fragments result from uptake of the antigen by a host cell, or synthesis of the antigen within the host cell, followed by cleavage of the antigen within the cell. Although it is probable that most successful vaccines elicit both T cell and B cell responses, current methods for evaluating test vaccines generally focus on antibody

production by B cells, and do not assess the ability of the test vaccine to elicit a T cell response.

Foreign proteins which are synthesized within the host cell or are taken up by the host cell via specific receptors are fragmented within the cytosol of the cell. One or
5 more of the resulting peptides can become associated with class I major histocompatibility molecules (MHC I), and the resulting complexes are then presented at the surface of the cell. These MHC I/peptide complexes are recognized by specific T cell receptors in certain CD8⁺ T cells, and the peptides so presented are referred to as CD8 epitopes.

10 A foreign protein can be taken up by a host cell nonspecifically via endocytosis and then fragmented into peptides in a cellular lysosomal or endosomal compartment. One or more of these peptides can then become associated with a class II major histocompatibility molecule (MHC II) to form a complex which is then presented at the surface of the host cell. These MHC II/peptide complexes are recognized by CD4⁺ T
15 cells expressing a specific receptor which recognizes the MHC II/peptide complex. These peptides are referred to as CD4 epitopes.

Peripheral T cells in the blood and organs of the immune system (e.g. spleen and lymph nodes) exist in a quiescent or resting state. Upon interaction of T cells with an MHC/epitope complex, the T cells proliferate and differentiate into activated cells
20 having a variety of functions. CD8⁺ T cells typically become cytotoxic upon activation and destroy antigen-presenting cells via direct contact. Activated CD4⁺ T cells provide a helper function to B cells, enabling B cells to differentiate into antibody-producing cells. Activated CD8⁺ T cells and CD4⁺ T cells release a variety of cytokines (lymphokines or interleukins), which can, for example, control differentiation of many
25 classes of lympholytic precursor cells.

In one embodiment, the invention provides a method for selecting one or more vaccine compositions from among a group of two or more vaccine compositions for *in vivo* assessment in one or more animals and/or human subjects. Each of the vaccine compositions comprises one or more antigens or one or more nucleic acid molecules encoding one or more antigens. The method comprises the steps of: (1) contacting
30 antigen presenting cells in culture with a vaccine composition selected from among said group of vaccine compositions, thereby, if one or more of the antigens or nucleic acid

00450473-003300

Chambers
14

molecules are taken up and processed by the antigen presenting cells, producing one or more processed antigens; (2) contacting the antigen presenting cells with T cells under conditions sufficient for the T cells to respond to one or more of the processed antigens; (3) determining whether the T cells respond to one or more of the processed antigens; 5 whereby if the T cells respond to one or more of the processed antigens, then the vaccine composition stimulates a T cell response; and (4) repeating steps (1), (2) and (3) with each vaccine composition in the group, thereby identifying vaccine compositions which stimulate a T cell response; and, if one or more of the vaccine compositions stimulates a T cell response, (5) selecting at least one vaccine composition which 10 stimulates a T cell response for assessment *in vivo*.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2

claim 15
5 nucleic acid molecules encoding one or more antigens to stimulate a protective T cell response. The method comprises the steps of: (1) contacting human antigen presenting cells in culture with the vaccine composition, thereby, if one or more of the antigens or nucleic acid molecules can be taken up and processed by the antigen presenting cells, producing one or more processed antigens; (2) contacting the antigen presenting cells with human T cells under conditions sufficient to produce a T cell response to one or more of the processed antigens, thereby producing a T cell response; (3) measuring the T cell response; and, if the T cell response is greater than a pre-selected value, (4) assessing the ability of the vaccine composition to stimulate a protective T cell response in one or more animals, human subjects or a combination thereof. The pre-selected value of the T cell response is, typically, chosen to represent a vaccine composition which is particularly active in stimulating a T cell response.

10
15 In another embodiment, the method of the invention comprises the steps of : (1) contacting human antigen presenting cells in culture with the vaccine composition, whereby, if one or more of the antigens are taken up and processed by the antigen presenting cells, said antigen or antigens are processed by the antigen presenting cells, thereby producing one or more processed antigens; (2) contacting the antigen presenting cells of step (1) with human T cell clones which are specific for an epitope within one or more of the antigens for a period of time sufficient for the human T cell clones to respond to one or more of the processed antigens; and (3) determining whether the human T cell clones respond to the processed antigen or antigens. If the T cell clones respond to the processed antigen or antigens, the method can, optionally, further include the step of assessing the vaccine composition in an animal or in a human.

20
25 A "processed antigen", as the term is used herein, refers to one or more epitopes derived from an antigen which are presented at the surface of an antigen presenting cell in combination with MHC I or MHC II.

30 The present method assesses the ability of a candidate vaccine composition to provide *in vitro* an antigen to antigen presenting cells in a manner which leads to processing and presentation of one or more T cell epitopes at the surface of the antigen presenting cells in combination with MHC I or MHC II. This *in vitro* determination provides an efficient screen for selecting compositions for more time-consuming *in vivo* testing in animals or in humans. This *in vivo* testing can be performed using methods

which are well known in the art. For example, the vaccine composition can be administered to an animal or a human, and the ability of the induced immune response, if any, to protect against subsequent challenge from the pathogen from which the antigen or antigens are derived can be determined. Alternatively, or in conjunction with
5 such a determination, the ability of the vaccine composition to induce *in vivo* the proliferation of T cells and/or antibodies which recognize one or more of the antigens can also be determined. Animals which can be used for *in vivo* testing include laboratory animals, domesticated animals and wild animals. Suitable examples include rodents, such as mice, hamsters, rats, guinea pigs and rabbits; primates, such as
10 monkeys and apes; and domestic animals, such as dogs, cats, horses, chickens, cows and pigs.

The antigen presenting cells are contacted with the vaccine composition in cell culture in a suitable culture medium, as is known in the art, and under suitable conditions, such as physiological pH, and at a temperature from about room temperature
15 to about physiological temperature, for a sufficient period of time for uptake and processing of the antigen by the antigen presenting cells. If the vaccine comprises a nucleic acid molecule, the antigen presenting cells are contacted with the vaccine composition for a sufficient amount of time for the antigen presenting cells to take up and express the nucleic acid molecule and process the resulting antigen. Generally, the
20 antigen presenting cells are contacted with the vaccine composition for a period of several hours, for example, from about 2 to about 12 hours. Following contact with the vaccine composition, the antigen presenting cells are contacted with the T cells for a sufficient period of time for activation of the T cells and generation of a T cell effector response. Generally, this process requires several hours, for example, from about 2 to
25 about 12 hours. Preferably, the APCs are contacted with the vaccine composition for a sufficient period time for antigen or nucleic acid molecule uptake, and then washed and placed in fresh media prior to addition of the T cells. Alternatively, the antigen presenting cells can be contacted with the vaccine composition and the T cells simultaneously or within a relatively short time interval. In this embodiment, the
30 antigen presenting cells are contacted with the vaccine composition and the T cells for a sufficient amount of time for antigen processing and generation of a T cell response. Typically, such a process requires from about 4 to about 24 hours.

The vaccine composition, preferably, comprises at least one antigen, or a nucleic acid encoding at least one antigen, which is a protein or a peptide which comprises one or more T cell epitopes, such as one or more CD8⁺ T cell epitopes, one or more CD4⁺ T cell epitopes or a combination thereof. Preferably, the T cells are specific for a particular epitope present within the antigen. More preferably, the T cells are T cell clones derived from a single precursor T cell. In a particularly preferred embodiment, the T cells are human T cell clones.

In one embodiment, the epitope is a CD4⁺ T cell epitope and the T cells are CD4⁺ T cells. As discussed above, the effector functions of CD4⁺ T cells include releasing cytokines and stimulating B cells to become antibody-producing cells. Thus, in this embodiment, the extent of the T cell response to the antigen presenting cells can be determined by measuring T cell proliferation, the production of one or more cytokines or the stimulation of antibody production by B cells. Greater levels of T cell proliferation, antibody production or cytokine production would be expected to correlate with greater immunogenicity and potential efficacy of the vaccine composition.

In another embodiment, the epitope is a CD8 epitope and the T cells are CD8⁺ T cells. As discussed above, the effector functions of CD8⁺ T cells include lysis of antigen presenting cells and release of cytokines. Therefore, the extent of CD8⁺ T cell response to the antigen presenting cells can be determined using an assay for cell lysis or by measuring the production of one or more cytokines. The CD8⁺ T cell response can also be measured by measuring the extent of release of one or more cytokines. In general, it is expected that greater cell lysis activity or cytokine release will correlate with greater immunogenicity.

The antigen presenting cells can be selected from among any suitable cells which are potentially capable of taking up the antigen, such as a natural, purified or recombinant protein, or a nucleic acid molecule encoding the antigen, and presenting a peptide epitope derived from the antigen at the cell surface in combination with MHC I or MHC II. For example, when the epitope is a CD4 epitope, cells expressing MHC II molecules can be used. Such cells include macrophages, dendritic cells and B cells. When the epitope is a CD8⁺ T cell epitope, the antigen presenting cells can be selected from among any cells which express MHC I. In preferred embodiments, the antigen-

presenting cells are professional antigen-presenting cells, such as macrophages, dendritic cells and B cells. The antigen presenting cells can be, for example, recombinant cells expressing heterologous MHC molecules. In a preferred embodiment, the antigen presenting cells are human cells. The antigen presenting cells
5 present the proper MHC molecules and are, preferably, at least partially HLA matched with the T cells. More preferably, the APCs are autologous cells, that is, cells derived from the same donor as the T cells.

In one embodiment, the T cells are clones which are specific for a particular epitope, and the vaccine composition includes at least one antigen which comprises the
10 epitope or at least one nucleic acid molecule encoding at least one antigen which comprises the epitope. In this embodiment, response of the epitope-specific T cell clones to antigen-presenting cells which have been contacted with the experimental vaccine composition indicates that the vaccine composition is able to effect the presentation of the epitope on the surface of the antigen-presenting cells in combination
15 with an MHC I or MHC II molecule.

Epitope-specific T cell clones can be generated using methods which are generally known in the art (see, for example, Fathman, *et al.*, in Paul, ed., *Fundamental Immunology*, second edition, Raven Press (1989), Chapter 30, the contents of which are hereby incorporated by reference in their entirety). The isolation of epitope-specific T
20 cell clones is based on T cell biology. Generally, an animal, such as a mouse, is immunized with a preparation of antigens (a bacterial lysate, or a purified protein) or is infected with a virus, such as a wild type virus or a recombinant virus containing heterologous genes encoding one or more proteins from a pathogenic microorganism, such as a virus. The animal is then sacrificed and the peripheral blood mononuclear
25 cells (PBMC: includes T cells, B cells, monocytes), spleen and lymph nodes are isolated. The isolated cells are then cultured in media containing a defined component of the original antigenic preparation, often a recombinant or purified protein, and the essential T cell growth factor interleukin-2 (IL-2). The only T cells which will proliferate are those which recognize MHC/epitope complex in which the epitope is
30 derived from the antigenic preparation. These cells become activated and proliferate while the unactivated cells begin to die. The cultures are maintained for several weeks, with the media containing antigen and IL-2 being periodically replaced. Eventually,

clusters of living and dividing cells (a T cell line) can be observed in some of the cultures.

manuscript
manuscript
5 The proliferating cells are generally not clonal at this point and are of limited use for assaying epitope specific T cell responses. The T cell line is, preferably, cloned through a process referred to as limiting dilution. In this method, PBMC are isolated from, for example, a mouse of the same strain as the original mouse used to isolate the T cell line. These cells, called antigen presenting cells, will serve as a source of MHC proteins and will present the MHC:peptide complex to the T cell line. The T cell line is diluted to a concentration of about 1 to 5 T cells/mL in a suspension of APCs that
10 contains the antigen of interest and IL-2. This suspension is then transferred into, for example, round or "v"-bottom 96 well microtitre plates, so that each well contains, on average, no more than 1 T cell. The cultures are maintained for several weeks and a clone can grow out of one or more cultures.

manuscript
15 The cells isolated by limiting dilution are the progeny of a single cell that expresses only one T cell receptor, and the clone is thus epitope-specific. However, in a situation in which the cloning procedure uses whole proteins or viruses, a single protein may contain many epitopes and the precise epitope will remain unknown. The epitope can be identified using a collection of overlapping synthetic peptides that span the entire amino acid sequence of the antigenic protein. These peptides can be used to
20 stimulate proliferation or cytokine secretion in a direct stimulation assay, or they may be used as competitive inhibitors to block activation of the T cell clone by the antigenic protein.

25 Human T cell clones can also be isolated. Generally, these clones are isolated from individuals who have had an infection, for example, influenza, HIV or Dengue, or have been exposed to antigens in nature or by injection and have T cells that specifically respond to those antigens. These antigens are called "recall antigens" and include tetanus toxoid and *Candida albicans* extract. Human T cell clones are isolated from the PBMC.

30 The T-cell response to APCs treated with the test vaccine composition can be determined using a variety of assays which are known in the art. Several examples are taught by Fathman, et al., *supra*. For example, T cell proliferation can be measured using methods known in the art. In one embodiment, the epitope-specific T cells are

435/6
mixed with irradiated antigen presenting cells and the test vaccine composition and cultured. The cells are cultured for a period of a few days to allow presentation of the epitope by the APCs and activation of the T cells. T cell proliferation is then assessed by monitoring the incorporation of ^3H -thymidine into newly synthesized DNA. The
5 APCs do not incorporate ^3H -thymidine because they have been irradiated. Alternative methods for assessing proliferation that do not use radioisotopes are also known.

10 T cell response can also be determined by determining if one or more cytokines is released by the T cells. For this assay, APCs and the test vaccine composition are mixed and cultured. Either simultaneously or after a period of time sufficient for uptake and processing of an antigen within the vaccine composition by the APCs, T cells are added to the culture. After a period of time sufficient to allow activation of the T cells, growth of the culture is stopped, for example, by freezing. Freezing the culture lyses the cells and releases cytokines that have not yet been secreted into the culture medium. The presence or absence of cytokine in the culture medium can then be determined
15 using known methods. Optionally, the amount of one or more cytokines in the culture medium can be determined. For example, cytokines in the culture supernatant and the cells can be measured using a bioassay, in which cell lines that proliferate only when stimulated with a particular cytokine (indicator cells) are cultured in media that is supplemented with an aliquot of the cytokine-containing culture media. The culture is maintained, typically, for 10-18 hours and ^3H -thymidine is added. After an additional
20 6-10 hours, new DNA synthesis is measured by determining the amount of ^3H incorporated into the cellular DNA. Any cytokine which is produced by the T cells upon activation can be measured. Examples of cytokines which can be determined include interferon- γ and interleukin-2.

25 In another embodiment, cytokine production is measured using an enzyme-linked immunosorbent assay (ELISA), for example, using reagents which are commercially available as kits. In this assay, an immobilized antibody is used to specifically capture a particular cytokine from the cytokine containing culture supernatant. Unbound proteins are washed away, and the amount of bound cytokine is
30 determined by binding a second, labeled, antibody to the captured cytokine. This assay is quantitative and more specific than bioassays. Alternatively, cytokine mRNA levels can be quantitated using the polymerase chain reaction. Cytokine production can also

435/6
Pundirich
ELISA

435/7.24
be determined by staining producer T cells with labeled antibodies specific for the cytokine.

In another embodiment, the T cells are CD8⁺ T cells and the response is measured by determining whether the T cells lyse the APCs which have been treated with the test vaccine composition. In one embodiment, the APCs are transformed peripheral blood lymphocyte cell lines (B-LCL) which have been incubated with ⁵¹CrO₄²⁻. The resulting ⁵¹Cr-labeled PBLs are thoroughly washed, incubated with the test vaccine composition and then exposed to the antigen-specific CD8⁺ T cells. After incubating for a sufficient period of time for epitope presentation by the B-LCLs and T cell activation, the extent of ⁵¹Cr release into the culture medium is determined. The amount of ⁵¹Cr released correlates with the extent of lysis of the B-LCLs.

The production of a T cell response can, generally, be determined by comparing the result achieved with the vaccine composition to a suitable control, as is known in the art. For example, in the ⁵¹Cr release assay discussed above, the amount of ⁵¹Cr released when the B-LCLs are treated with the CD8⁺ T cells can be compared to the amount released when the B-LCLs are treated with vehicle alone, referred to as the background release. Significantly (measurably) greater ⁵¹Cr release in the presence of the T cells is indicative of a T cell response. In the cytokine production assay, cytokine production by the T cells in the presence of APCs treated with the vaccine composition can be compared to cytokine production by the T cells in the absence of APCs, or in the presence of untreated APCs. Greater cytokine production in the presence of treated APCs is indicative of a T cell response.

The test vaccine composition comprises one or more antigens or one or more nucleic acid molecules which encode one or more antigens. The vaccine composition can be any of the types of vaccine compositions which are known in the art. For example, the vaccine composition can comprise an attenuated pathogen, such as a weakened bacterial strain or virus, or a killed pathogen, such as a killed bacterial strain or a killed virus. The vaccine composition can also comprise a portion of a pathogen, for example, a viral coat or bacterial membrane. In another embodiment, the vaccine composition comprises one or more proteins derived from a pathogen, for example, a protein which has been purified or partially purified from the pathogen, or a recombinant protein produced by a recombinant organism which expresses a gene

derived from the pathogen which encodes the protein. Examples of suitable host organisms for the production of recombinant peptides and proteins are known in the art and include *E. coli*. The vaccine composition can also include one or more fragments of a protein or proteins derived from pathogen. Such protein fragments include peptides which are synthesized or recombinantly produced.

In another embodiment, the test vaccine composition includes one or more proteins, or fragments thereof, which are produced by a particular type of tumor cell. Preferably, the protein is unique to the tumor cell, i.e., not present in or on healthy cells, or is expressed in greater quantity by the tumor cell than by healthy cells. The protein can be, for example, a protein found on the surface of the tumor cell. The protein(s) can be derived from the tumor cells, for example, isolated and purified or partially purified from cultured tumor cells. The tumor cell protein(s), or a fragment or fragments thereof, can also be produced recombinantly.

In another embodiment, the vaccine composition comprises a nucleic acid molecule which encodes a protein or a fragment thereof, derived from a pathogen or a tumor cell as discussed above. For example, the vaccine composition can comprise so-called "naked DNA". The nucleic acid molecule can also be contained within a suitable vector, such as a recombinant virus, such as vaccinia virus, adenovirus, orf virus, fowlpox virus, herpes virus, varicella virus, papilloma virus, SV40, retroviruses, baculovirus and poliomyelitis virus. The vector can also be a bacterium, such as salmonella, BCG or *E. coli*. The nucleic acid can also be present in a liposome or another suitable vector, such as are known in the art.

As discussed above, the present invention enables the rapid assessment and comparison of a large number of potential vaccine compositions. For any given disease or pathogen, for example, a variety of antigens can be assessed. For example, a set of vaccine compositions which each include different antigens or portions of antigens from a particular pathogen can be compared. Further, for a given antigen, set of antigens, or nucleic acid molecule encoding such antigen(s), a variety of formulations can be assessed. For example, a set of vaccine compositions including the same antigen or antigens, but different vectors, adjuvants, concentrations, vehicles or excipients can be compared to determine the conditions necessary for optimal efficacy.

The invention will now be further and specifically described in the following examples.

EXAMPLES

Materials and Methods

5 Viruses

Influenza A viruses A/Puerto Rico/8/34 (H1N1) and A/Japan/305/57 (H2N2) were obtained from the Division of Virology, Bureau of Biologics, Food and Drug Administration, Bethesda, Md. A/Johannesburg/94 (H3N2) was obtained from David Burt (Pasteur Merieux Connaught, Toronto, Ontario, Canada). Influenza A viruses were
10 propagated in 10-day-old, embryonated chicken eggs. Infected allantoic fluids were harvested 2 days after infection, aliquoted, and stored at -80°C until use. Recombinant vaccinia viruses containing the genes coding for influenza A viral proteins HA, NA, M1, M2, PB1, PB2, PA, NS1, and NS2 and the nucleoprotein (NP) were obtained from B. Moss. Each of these was derived from the A/PR/8/34 influenza A virus strain,
15 except for NS1, which was derived from A/Udorn/72. They were constructed and propagated as previously described (Smith *et al.*, *Virology* **160**: 336-345 (1987)). A recombinant vaccinia virus which expressed segmented portions of the NP was obtained from J. Bennink and L. Eisenlohr.

Human PBMC

20 PBMC specimens were obtained from normal, healthy donors. Most of the donors whose PBMC were tested had convincing evidence of influenza A virus-specific CTL activity in bulk culture. PBMC were purified by Ficoll-Hypaque density gradient centrifugation (A. Boyam, *Scand. J. Clin. Lab. Invest.* **21**: 77-89 (1968)). Cells were resuspended at 2×10^7 /mL in RPMI 1640 with 20% fetal bovine serum (FBS) (Sigma)
25 and 10% dimethyl sulfoxide and cryopreserved until use. The HLA alleles of donor 1 were A2.1, A11, B18, B27, Cw1, Cw7, DR1, DQw1, DQw3, DRw52, and DRw53. HLA typing was performed in the HLA typing laboratory at the University of Massachusetts Medical Center.

Bulk cultures of PBMC

Responder PBMC were suspended at $10^6/\text{mL}$ in AIM-V medium (Gibco BRL, Grand Island, N.Y.) containing 10% human AB serum (NABI, Boca Raton, Fla.), penicillin-streptomycin, glutamine, and HEPES in a 70-mL flask (Falcon). Stimulators
5 were infected with A/PR/8/34 at a multiplicity of infection (MOI) of 15 for 1.5 h at 37°C in 1 mL of phosphate-buffered saline containing 0.1% bovine serum albumin and then added to responders in a flask at a stimulator-responder ratio of 1:10. On day 7 of culture, cells were either cloned by limiting dilution as described below or restimulated with gamma-irradiated (3,000 rads) autologous PBMC infected with A/PR/8/34 at an
10 MOI of 15 for 1.5 h in 1 mL of phosphate-buffered saline containing 0.1% bovine serum albumin, added at a stimulator-responder ration of 1:10 in fresh medium containing 10% human AB serum and 20 U of interleukin-2 (IL-2) (Collaborative Biomedical Products, Bedford, Mass.). Restimulated cells were either cloned by limiting dilution or assayed for cytolytic activity 7 days later.

15 CTL clones

Influenza virus-specific CTL clones were established by using a limiting-dilution technique as previously described (Kurane *et al.*, *J. Exp. Med.* **170**: 763-775 (1989)). PBMC which had been stimulated in bulk culture for 7 or 14 days were collected and plated at a concentration of 3, 10, or 30 cells per well in 96-well round-
20 bottom microtiter plates in 100 μL of AIM-V medium containing 10% FBS, 20 U of IL-2, a 1:1,000 dilution of anti-CD8 monoclonal antibody 12F6 (obtained from Johnson Wong), and 10^5 gamma-irradiated allogeneic PBMC/well. On day 7, 50 μL of fresh medium with FBS (Sigma Immunochemicals, St. Louis, Mo.) and IL-2 were added, and on day 14, fresh medium with 10^5 gamma-irradiated allogeneic PBMC/well and a
25 1:1,000 dilution of the anti-CD8 monoclonal antibody were added. Growing cells were assayed for cytolytic activity on days 21 and 28. Cells from wells with influenza A virus-specific cytolytic activity were expanded to 48-well plates.

Preparation of target cells

Autologous lymphoblastoid cell lines (B-LCLs) were established by culturing
30 with Epstein-Barr virus in 24-well plates as previously described (Green *et al.*, *J. Virol.*

67: 5962-5967 (1993)). B-LCL were infected with recombinant vaccinia viruses at an MOI of 20:1 for 1.5 h at 37°C. The cells were then diluted in 1 mL of medium and further incubated for 12 to 16 h. Other B-LCL were infected with A/PR/8/34, A/Japan/305/57, or A/Johannesburg/94 in 1 mL of medium for 12 to 16 h. These
5 infected target cells were labeled with 0.25 mCi of ^{51}Cr for 60 min at 37°C. After four washes, the target cells were counted and diluted to $2 \times 10^4/\text{mL}$ for use in the cytotoxicity assay. The partially HLA-matched allogeneic target cells used in the assays were B-LCL produced in our laboratory from the HLA-typed PBMC of unrelated donors or were obtained from the National Institute of General Medical Sciences
10 Human Genetic Mutant Cell Repository or the American Society for Histocompatibility and Immunogenetics Cell Bank and Repository.

Cytotoxicity assays

Cytotoxicity assays were performed with 96-well round-bottom plates. Effector cells in 100 μL of RPMI 1640 medium containing 10% FBS were added to 2×10^5 ^{51}Cr -
15 labeled target cells in 100 μL at an effector-to-target (E-T) ratio of 10:1. Plates were centrifuged at $200 \times g$ for 5 min and incubated for 4 to 5 h at 37°C. Supernatant fluids were harvested by using the supernatant collection system (Skatron Instruments, Sterling, Va.), and ^{51}Cr content was measured in a gamma counter. Percent specific ^{51}Cr release was calculated with the following formula: $(\text{cpm experimental release} - \text{cpm spontaneous release}) / (\text{cpm maximum release} - \text{cpm spontaneous release}) \times 100$. All
20 assays were performed in triplicate, and the results were calculated from the average of the triplicate wells.

Example 1 *In Vitro* Evaluation of Influenza Vaccine Compositions

This Example was designed to evaluate whether proper formulation of an
25 influenza virus comprising a formalin-inactivated detergent-disrupted virus can lead to a CD8^+ cytotoxic T cell response.

The HA-Specific CD8^+ cytotoxic T cell clone described in Example 1 was incubated with autologous B-LCL cells which had been treated with one of the following vaccine formulations:

1. live influenza virus (H1N1, A/PR/8/34, A/Texas/9I)
2. iscomatrix alone
3. formalin-inactivated A/Texas/H1N1 virus; and
4. formalin-inactivated A/Texas/H1N1 virus formulated with Iscoms.

5 The results are shown in Table 1, which provides cytolysis as a per cent of total APCs and the background ^{51}Cr release. The data show that the CD8^+ clone recognized APC infected with live flu virus or a recombinant vaccinia virus. This CD8^+ CTL clone, however, did not lyse APC pulsed with the inactivated A/Texas/H1N1 virus unless it was formulated with an adjuvant carrier. Formulation with Iscoms enabled
10 processing of the vaccine for CD8^+ CTL recognition.

Table 1: Per cent lysis of B-LCLs treated with vaccine formulations by HA-specific cytotoxic T cell clone

	PR/8 virus	A/Tx virus	Iscomatrix	A/Tx vaccine	A/Tx Flu - Iscoms
CD8^+ clone	36.1%	19.5%	-3.5%	-2.2%	86.9%
15 min/max	12.8%	12.1%	17.4%	41.3%	44.6%

Example 2 *In Vitro* evaluation of recombinant protein vaccine formulations

A human CD8^+ cytotoxic T cell clone that recognizes amino acids 122-130 of the Influenza A NS1 protein is disclosed in U.S. Patent No. 5,766,601, the teachings of which are incorporated herein by reference in their entirety. This T cell clone was
20 incubated with autologous ^{51}Cr -labeled B-LCL treated with (1) a synthetic peptide based on NS1 (aa 122-130); (2) recombinant NS1 protein; or (3) recombinant NS1 protein formulated as an Iscom. Controls were also established using uninfected B-LCL and B-LCL incubated with Iscoms only.

The results of this study are presented in Table 2. The CD8⁺ cytotoxic T cell clone lysed APC that were treated with the recombinant NS1 aa 122-130 peptide and the recombinant NS1 protein/Iscom formulations. The CTL clone did not lyse APC treated with the recombinant protein alone or either of the control cells.

5 Table 2: Per cent lysis of autologous B-LCL treated with the indicated formulations by autologous NS1 aa 122-130 specific CD8⁺ cytotoxic T cell clone

	uninfected	NS1 aa 122-130 a	NS1	NS1 protein Iscom b,c	Iscom d
CD8 ⁺ clone	-6.2%	88.4	-0.7%	19.6%	-11.0
10 min/max	17.8%	19.6	-17.3%	29.9%	21.3%

a Peptide used at 25 µg/mL; b Protein used at 35 µg/mL;

c Saponin used at 100 µg/mL; d Saponin used at 70 µg/mL

Example 3 *In vitro* evaluation of HIV-1 vaccine compositions

A human HIV-1 specific CD8⁺ cytotoxic T cell clone was prepared as described
 15 by Littaua *et al.*, *J. Virol.* **65** : 4051-4056 (1991), the teachings of which are hereby incorporated by reference in their entirety. The ability of this clone to recognize autologous APCs pulsed with (1) recombinant HIV-1 p24 protein alone; (2) HIV-1 p24 in an Iscom formulation; or (3) a recombinant vaccinia virus containing the HIV-1 p24 gene was determined.

20 The results are presented in Table 3, which shows that APCs treated with the recombinant vaccinia virus are significantly lysed by the T cell clone. APCs treated with the p24/Iscoms complex are also recognized by the T cell clone, but to a lesser extent. The T cell clone did not recognize B-LCLs pulsed with the recombinant p24 protein alone.

Table 3: Lysis of B-LCL treated with indicated formulations by human HIV-1 specific CD8⁺ cytotoxic T cell clone

Vac/p24	Iscoms + p24 at			Iscom	Media
	100 µg	50 µg	25 µg	100 µg	
37.5	11.4	20.2	-0.1	-2.6	0.3

5 Example 5 *In vivo* evaluation of influenza virus compositions

Fifty-five healthy adults from 18 to 45 years old were enrolled in 5 study groups of 11 participants each: 1. Fluzone; 2. Flu-Iscom (75 µg); 3. Flu-Iscom (50 µg); 4. Flu-Iscomatrix (75 µg); and 5. Flu-Iscomatrix (50 µg). Cytotoxic T cell activity in the peripheral blood lymphocytes of the subjects was determined on days 0, 14 and 56 following a single immunization with trivalent vaccine. Peripheral blood lymphocytes at each time point for each subject were tested in the same assay for killing of virus-infected autologous target cells (Epstein-Barr virus transformed B cells) at various E:T ratios (90, 30 and 10). Responders were those subjects which showed a significant increase in killing of greater than 5% compared with the percent net lysis at time 0 at two or more effector:target ratios. The results are presented in Table 4.

Table 4

Virus Strain	Fluzone alone 50/45	Flu-Iscom 75/45	Flu-Iscom 50/45	Flu MTRX 75/45	Flu MTRX 50/45
H1	0/11	7/11	5/11	1/11	5/11
H3	1/11	3/11	7/11	3/11	5/11
B	5/11	6/11	5/11	6/11	7/11

These results show that the effect of the adjuvant was significant for virus strains H1 and H3, while results were similar for strain B in the presence and absence of an adjuvant. Figures 1A and 1B also present data illustrating the increase in net lysis

compared to day 0 for strain A/Texas at days 14 and 56, respectively. Similar data for strain A/Johannesburg are illustrated in Figures 2A and 2B.

EQUIVALENTS

While this invention has been particularly shown and described with references
5 to preferred embodiments thereof, it will be understood by those skilled in the art that
various changes in form and details may be made therein without departing from the
spirit and scope of the invention as defined by the appended claims. Those skilled in
the art will recognize or be able to ascertain using no more than routine
experimentation, many equivalents to the specific embodiments of the invention
10 described specifically herein. Such equivalents are intended to be encompassed in the
scope of the claims.